

in patients with valvular aortic stenosis. Second, the stepped and sequential LVPO induced in the present study is unlikely to completely represent the duration, pattern, and magnitude that occur in clinical forms of LVPO. Nevertheless, this model of LVPO demonstrated significant phenotypic similarities to clinical observations of LVPO in patients in terms of myocardial ECM remodeling and LV diastolic dysfunction. The model system and paradigms identified in this article may provide a means to identify the mechanisms that give rise to pathologic ECM remodeling and the potential point of irreversible ECM remodeling, and thereby improve the timing and treatment approaches for elimination of LVPO stimulus.

References

- Monrad ES, Hess OM, Murakami T, et al. Time course of regression of left ventricular hypertrophy after aortic valve replacement. *Circulation*. 1988;77:1345-55.
- Villari B, Vassalli G, Monrad ES, et al. Normalization of diastolic dysfunction in aortic stenosis late after valve replacement. *Circulation*. 1995;91:2353-8.
- Krayenbuehl HP, Hess OM, Monrad ES, et al. Left ventricular myocardial structure in aortic valve disease before, intermediate, and late after aortic valve replacement. *Circulation*. 1989;79:744-55.
- Hess OM, Ritter M, Schneider J, et al. Diastolic stiffness and myocardial structure in aortic valve disease before and after valve replacement. *Circulation*. 1984;69:855-65.
- Azevedo CF, Nigri M, Higuchi ML, et al. Prognostic significance of myocardial fibrosis quantification by histopathology and magnetic resonance imaging in patients with severe aortic valve disease. *J Am Coll Cardiol*. 2010;56:278-87.
- Villar AV, Llano M, Cobo M, et al. Gender differences of echocardiographic and gene expression patterns in human pressure overload left ventricular hypertrophy. *J Mol Cell Cardiol*. 2009;46:526-35.
- Kuwahara F, Kai H, Tokuda K, et al. Transforming growth factor-beta function blocking prevents myocardial fibrosis and diastolic dysfunction in pressure-overloaded rats. *Circulation*. 2002;106:130-5.
- Moorjani N, Ahmad M, Catarino P, et al. Activation of apoptotic caspase cascade during the transition to pressure overload-induced heart failure. *J Am Coll Cardiol*. 2006;48:1451-8.
- Schubert A, Walther T, Falk V, et al. Extracellular matrix gene expression correlates to left ventricular mass index after surgical induction of left ventricular hypertrophy. *Basic Res Cardiol*. 2001;96:381-7.
- Nagatomo Y, Carabello BA, Coker ML, et al. Differential effects of pressure or volume overload on myocardial MMP levels and inhibitory control. *Am J Physiol Heart Circ Physiol*. 2000;278:H151-61.
- Walther T, Schubert A, Falk V, et al. Regression of left ventricular hypertrophy after surgical therapy for aortic stenosis is associated with changes in extracellular matrix gene expression. *Circulation*. 2001;104(Suppl 1):I-54-8.
- Deardorff R, Spinale FG. Cytokines and matrix metalloproteinases as potential biomarkers in chronic heart failure. *Biomark Med*. 2009;3:513-23.
- Deschamps AM, Spinale FG. Pathways of matrix metalloproteinase induction in heart failure: bioactive molecules and transcriptional regulation. *Cardiovasc Res*. 2006;69:666-76.
- Nimmi ME. Fibrillar collagens: their biosynthesis, molecular structure, and mode of assembly. In: Zern MA, Reid LM, eds. *Extracellular Matrix*. New York: Marcel Dekker; 1993:121-48.
- López B, González A, Hermida N, et al. Role of lysyl oxidase in myocardial fibrosis: from basic science to clinical aspects. *Am J Physiol Heart Circ Physiol*. 2010;299:H1-9.
- Yarbrough WM, Mukherjee R, Brins TA, et al. Matrix metalloproteinase inhibition modifies left ventricular remodeling after myocardial infarction in pigs. *J Thorac Cardiovasc Surg*. 2003;125:602-10.
- Zile MR, DeSantis SM, Baicu CF, et al. Plasma biomarkers that reflect determinants of matrix composition identify the presence of left ventricular hypertrophy and diastolic heart failure. *Circ Heart Fail*. 2011;4:246-56.
- Spinale FG, Coker ML, Thomas CV, et al. Time-dependent changes in matrix metalloproteinase activity and expression during the progression of congestive heart failure: relation to ventricular and myocyte function. *Circ Res*. 1998;82:482-95.
- Lindsey ML, Goshorn DK, Squires CE, et al. Age-dependent changes in myocardial matrix metalloproteinase/tissue inhibitor of metalloproteinase profiles and fibroblast function. *Cardiovasc Res*. 2005;66:410-9.
- Marotta M, Martino G. Sensitive spectrophotometric method for the quantitative estimation of collagen. *Anal Biochem*. 1985;150:86-90.
- Walsh BJ, Thornton SC, Penny R, Breit SN. Microplate reader-based quantitation of collagens. *Anal Biochem*. 1992;203:187-90.
- Dahl JS, Videbaek L, Poulsen MK, et al. Effect of candesartan treatment on left ventricular remodeling after aortic valve replacement for aortic stenosis. *Am J Cardiol*. 2010;106:713-9.
- Monrad ES, Hess OM, Murakami T, et al. Abnormal exercise hemodynamics in patients with normal systolic function late after aortic valve replacement. *Circulation*. 1988;77:613-24.
- Ahmed SH, Clark LL, Pennington WR, et al. Matrix metalloproteinases/tissue inhibitors of metalloproteinases: relationship between changes in proteolytic determinants of matrix composition and structural, functional, and clinical manifestations of hypertensive heart disease. *Circulation*. 2006;113:2089-96.
- Heymans S, Schroen B, Vermeersch P, et al. Increased cardiac expression of tissue inhibitor of metalloproteinase-1 and tissue inhibitor of metalloproteinase-2 is related to cardiac fibrosis and dysfunction in the chronic pressure-overloaded human heart. *Circulation*. 2005;112:1136-44.
- Foronjy RF, Sun J, Lemaitre V, et al. Transgenic expression of matrix metalloproteinase-1 inhibits myocardial fibrosis and prevents the transition to heart failure in a pressure overload mouse model. *Hypertens Res*. 2008;31:725-35.
- Burke B. The role of matrix metalloproteinase 7 in innate immunity. *Immunobiology*. 2004;209:51-6.
- Spinale FG, Mukherjee R, Zavadzka JA, et al. Cardiac restricted overexpression of membrane type-1 matrix metalloproteinase causes adverse myocardial remodeling following myocardial infarction. *J Biol Chem*. 2010;285:30316-27.
- Tummalapalli CM, Heath BJ, Tyagi SC. Tissue inhibitor of metalloproteinase-4 instigates apoptosis in transformed cardiac fibroblasts. *J Cell Biochem*. 2001;80:512-21.
- Stetler-Stevenson WG. The tumor microenvironment: regulation by MMP-independent effects of tissue inhibitor of metalloproteinases-2. *Cancer Metastasis Rev*. 2008;27:57-66.
- Lovelock JD, Baker AH, Gao F, et al. Heterogeneous effects of tissue inhibitors of matrix metalloproteinases on cardiac fibroblasts. *Am J Physiol Heart Circ Physiol*. 2005;288:H461-8.
- Yarbrough WM, Mukherjee R, Ikonomidis JS, Zile MR, Spinale FG. Myocardial remodeling with aortic stenosis and after aortic valve replacement: mechanisms and future prognostic implications. *J Thorac Cardiovasc Surg*. 2011 Jul 13 [Epub ahead of print].

Discussion

Dr Joseph Schmoker (Burlington, Vt). I have no disclosures. My questions pertain to the miR-1 and 133 that you studied. In the mouse model, the transverse aortic constriction (TAC) model, both are actually antihypertrophic and overexpression of miR-133 is associated with down-regulation of collagen in cultured fibroblasts. So I'm curious how you interpret this relative to the opposite findings in your large animal model. In addition, did you perform microarray analysis on myocardium to broaden your miR phenotype?

Dr Yarbrough. We have just begun our initial assessments with respect to miRs. We anticipate performing microarray analysis to broaden the miR phenotype. I agree these initial preliminary miR data are at odds with the relatively small amount of data that exist in the literature with respect to this topic.

At this time, we believe that miR-133 in particular may inhibit collagen 1A1 and 3A1. Our theory is that with up-regulation of miR-133, there is inhibition of collagen 1A1 and 3A1 expression. That would account for the lack of increased collagen expression

noted in the study. In summary, we believe the increased collagen content observed in the study was secondary to increased collagen stability and resistance to degradation as opposed to increased collagen expression. With prolongation of the animal model, we may see a reduction in miR-133 and miR-1 levels that will then “release” some of the inhibition on the collagen 1A1 and 3A1 genes. That may then lead to a robust increase in collagen expression, counter to what we’re seeing at this point in time at 1 month.

Dr Schmoker. My final question is related to how the control animals were handled at the terminal data collection. My understanding from the article is that the controls were not violated

from the standpoint of manipulation of the heart during final data collection. Knowing that both miR-1 and 133 can be induced with myocardial ischemia, could subendocardial ischemia incurred from surgical handling and manipulation during terminal data collection be responsible for the differences in miR expression in the experimental group relative to controls?

Dr Yarbrough. Perhaps, but I don’t believe that is likely. We performed microsphere studies on both the control and the instrumented animals and showed no difference in perfusion with respect to microsphere data collected from the endocardium, myocardium, or epicardium. Thus, I don’t think submyocardial ischemia was likely.